

routine immunisation schedule from January 2020

| Age due | Diseases protected against | Vaccine given and trade name | | Usual site |
|------------------------------------|--|---|-------------------------------------|-----------------|
| 12 weeks old | Diphtheria, tetanus, pertussis (whooping cough), polio, <i>Haemophilus influenzae</i> type b (Hib) and hepatitis B | DTaP/IPV/Hib/HepB | Infanrix hexa | Thigh |
| | Meningococcal group B (MenB) | MenB | Bexsero | Left thigh |
| | Rotavirus gastroenteritis | Rotavirus | Rotarix | By mouth |
| 16 weeks old | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B | DTaP/IPV/Hib/HepB | Infanrix hexa | Thigh |
| | Pneumococcal (13 serotypes) | Pneumococcal conjugate vaccine (PCV) | Prevenar 13 | Thigh |
| | Rotavirus | Rotavirus | Rotarix | By mouth |
| 20 weeks old | Diphtheria, tetanus, pertussis, polio, Hib and hepatitis B | DTaP/IPV/Hib/HepB | Infanrix hexa | Thigh |
| | MenB | MenB | Bexsero | Left thigh |
| 1 year old (at birthday) | Hib and MenC | Hib/MenC | Menitorix | Upper arm/thigh |
| | Pneumococcal | PCV booster | Prevenar 13 | Upper arm/thigh |
| | Measles, mumps and rubella (German measles) | MMR | MMR VaxPRO ² or Priorix | Upper arm/thigh |
| | MenB | MenB booster | Bexsero | Left thigh |
| | Influenza (each year from September) | Live attenuated influenza vaccine LAIV ^{2,3} | Fluenz Tetra ^{2,3} | Both nostrils |
| 1 year 4 months old or soon | Diphtheria, tetanus, pertussis and polio | DTaP/IPV | Repevax or Boostrix-IPV | Upper arm |
| | Measles, mumps and rubella | MMR (check first dose given) | MMR VaxPRO ² or Priorix | Upper arm |
| 13 years old (girls aged thirteen) | Cancers caused by human papillomavirus (HPV) types 16 and 18 (and genital warts caused by types 6 and 11) | HPV (two doses 6-24 months apart) | Gardasil | Upper arm |
| 1 year 9 months old (at 9) | Tetanus, diphtheria and polio | Td/IPV (check MMR status) | Revaxis | Upper arm |
| | Meningococcal groups A, C, W and Y disease | MenACWY | Nimenrix or Menveo | Upper arm |
| 2 years old | Pneumococcal (23 serotypes) | Pneumococcal Polysaccharide Vaccine (PPV) | Pneumococcal Polysaccharide Vaccine | Upper arm |
| 5 years of age | Influenza (each year from September) | Inactivated influenza vaccine | Multiple | Upper arm |
| 50 years old | Shingles | Shingles | Zostavax ² | Upper arm |

See the green book chapter 19 or visit www.gov.uk/government/publications/influenza-the-green-book-chapter-19 or www.gov.uk/government/publications/influenza-the-green-book-chapter-19/conditions/vaccinations/child-flu-vaccine/

contains porcine gelatine.

If an inactivated influenza vaccine is contraindicated and child is in a clinical risk group, use inactivated flu vaccine.

For more information on vaccine supply information for the routine immunisation schedule please visit www.immform.dh.gov.uk
 For more information on vaccine supply information for all other vaccine supply information.

